

Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics

www.renhyd.org

RESEARCH ARTICLE

Effects of Plant Extract and Dietary Supplements on Tinnitus Perception in Adults: A Systematic Review and Meta-Analysis

Yordanis Enriquez Canto^{a,*}, Giovani Díaz Gervasi^a, Luis Menacho Alvirio^b, Rafael Pizarro-Mena^c

^a Facultad de Ciencias de la Salud, Universidad Católica Sedes Sapientiae, Lima, Perú.

^b Instituto de Medicina Tropical Alexander Von Humboldt, Universidad Peruana Cayetano Heredia, Lima, Perú.

^c Facultad de Ciencias de la Rehabilitación y Calidad de Vida, Universidad San Sebastián, Sede Los Leones, Santiago, Chile.

*yenriquez@ucss.edu.pe

Assigned Editor: Malak Kouiti. Universidad Hassan Primero de Settat, Settat, Marruecos.

Received: 09/12/2024; Accepted: 01/16/2025; Published: 03/14/2025.

KEYWORDS

Tinnitus;
Dietary Supplements;
Melatonin;
Plant Extracts;
Randomized
Controlled Trials.

Effects of Plant Extract and Dietary Supplements on Tinnitus Perception in Adults: A Systematic Review and Meta-Analysis

ABSTRACT

Introduction: Tinnitus, the perception of sound without an external source, affects millions worldwide and significantly impacts the health-related quality of life (HRQoL). Despite various management strategies, no universally effective treatment exists. While tinnitus sufferers frequently use dietary supplements, their efficacy remains unclear. This systematic review and meta-analysis evaluated the effectiveness of plant extracts and dietary supplements in treating tinnitus symptoms.

Methodology: We conducted a systematic search in PubMed for randomized controlled trials (RCTs) and crossover trials comparing plant extracts and dietary supplements with placebo or control groups in adults with tinnitus. Selection criteria included patients with subjective tinnitus and studies examining plant extracts or dietary supplements. Using a random-effects model, we estimated standardized mean differences and odds ratios with 95% confidence intervals (95% CIs). Study quality was assessed using the Revised Cochrane risk-of-bias tool for randomized trials. Protocol registered on PROSPERO (CRD42023446824).

Results: Twenty-one studies involving 1,759 participants evaluated four plant extracts and six dietary supplements. Five trials investigated plant extracts (*Euterpe oleracea*, traditional herbal medicine Hangekobokuto, Neurotec, and caffeine), while sixteen studies examined dietary supplements (melatonin, zinc, alpha-lipoic acid, manganese, vitamin B complex, and nicotinamide). Meta-analysis revealed no significant efficacy for caffeine in reducing tinnitus impact on HRQoL (SMD=0.07; 95%CI: -0.73 to 0.86; $I^2=0.0\%$). However, melatonin showed a modest reduction in tinnitus loudness compared to placebo (OR=2.52; 95%CI: 1.33-4.80; $k=2$, 168 participants).

Conclusions: Among the studied interventions, only melatonin showed modest efficacy in reducing tinnitus loudness, though evidence remains uncertain due to methodological limitations. Further high-quality RCTs with larger samples, standardized outcome measures, and long-term follow-up are necessary to confirm these findings and explore melatonin's potential role in tinnitus management.



➤ **Efectos de los extractos de plantas y los suplementos dietéticos en la percepción del tinnitus en adultos:
Una revisión sistemática y metaanálisis**

PALABRAS CLAVE

Tinnitus;
Suplementos
Dietéticos;
Melatonina;
Extractos de Plantas;
Ensayos Controlados
Aleatorizados.

RESUMEN

Introducción: El tinnitus, la percepción de sonido sin una fuente externa, afecta a millones de personas en todo el mundo y tiene un impacto significativo en la calidad de vida relacionada con la salud (CVRS). A pesar de las diversas estrategias de manejo, no existe un tratamiento universalmente efectivo. Si bien los afectados por tinnitus utilizan frecuentemente suplementos dietéticos, su eficacia sigue siendo incierta. Esta revisión sistemática y metaanálisis evaluó la efectividad de los extractos de plantas y los suplementos dietéticos en el tratamiento de los síntomas del tinnitus.

Metodología: Realizamos una búsqueda sistemática de ensayos controlados aleatorios (ECA) o ensayos cruzados que compararan extractos de plantas y suplementos dietéticos con grupos de placebo o control en adultos con tinnitus. Los criterios de selección incluyeron pacientes con tinnitus subjetivo y estudios que examinaran extractos de plantas o suplementos dietéticos. Utilizando un modelo de efectos aleatorios, estimamos las diferencias de medias estandarizadas y las razones de momios con intervalos de confianza del 95% (IC del 95%). La calidad de los estudios se evaluó utilizando la herramienta de riesgo de sesgo revisada de Cochrane para ensayos aleatorios. Protocolo registrado en PROSPERO (CRD42023446824).

Resultados: 21 estudios involucraron a 1.759 participantes y evaluaron cuatro extractos de plantas y 6 suplementos dietéticos. 5 ensayos investigaron extractos de plantas (*Euterpe oleracea*, medicina herbal tradicional Hangekobokuto, Neurotec y cafeína), mientras que 16 examinaron suplementos dietéticos (melatonina, zinc, ácido alfa-lipoico, manganeso, complejo de vitamina B y nicotinamida). El metaanálisis reveló que no hubo una eficacia significativa de la cafeína en la reducción del impacto del tinnitus en la CVRS (DME=0,07; IC del 95%: -0,73 a 0,86; $I^2=0,0\%$). La melatonina mostró una reducción modesta en la intensidad del tinnitus vs. placebo (OR=2,52; IC del 95%: 1,33-4,80; $k=2$, 168).

Conclusiones: Solo la melatonina mostró una eficacia modesta en la reducción del tinnitus, aunque la evidencia sigue siendo incierta debido a limitaciones metodológicas. Se necesitan más investigaciones de calidad con seguimiento a largo plazo que confirmen estas observaciones.

**KEY
MESSAGES**

1. Melatonin shows modest potential in reducing tinnitus loudness, but evidence remains uncertain due to small sample sizes ($n=46-122$), high dropout rates ($>20\%$) and variable study quality.
2. Evaluated interventions including *Euterpe oleracea* extract, traditional herbal medicine (Hangekobokuto), commercial formulation (Neurotec), caffeine, and dietary supplements showed no significant effect on tinnitus symptoms or HRQoL.
3. High heterogeneity in outcome measures limits evidence synthesis and impedes clinicians' ability to make informed treatment decisions, highlighting the need for standardized assessment tools to guide treatment selection in clinical practice.
4. Clinical trials should define meaningful improvement as ≥ 20 -point reduction in THI/TFI scores, include both subjective (VAS for loudness/annoyance) and objective measures, and report adverse events systematically to enable evidence-based recommendations for dietary supplements or plant-based interventions.

CITATION

Enriquez Canto Y, Díaz Gervasi G, Menacho Alvirio L, Pizarro-Mena R. Effects of Plant Extract and Dietary Supplements on Tinnitus Perception in Adults: A Systematic Review and Meta-Analysis. Rev Esp Nutr Hum Diet. 2025; 29(1): 7-20. doi: <https://doi.org/10.14306/renhyd.29.1.2308>

INTRODUCTION

Tinnitus, the perception of sound without an external source, is a global burden affecting between 4.1% and 37.2% of adults^{1,2}. It manifests as ringing or hissing in the ears or head, resulting from the brain's misinterpretation of abnormal auditory activity¹. While both objective (sound heard by patient and examiner) and subjective (sound heard only by patient) forms exist, subjective tinnitus is more common and occurs without organic or pathological lesions^{3,4}. Its prevalence increases with age, affects both sexes equally, and persistently bothers 3-31% of sufferers³. The condition significantly impacts quality of life, causing difficulties with concentration, sleep, and communication, often triggering anxiety and depression⁵. Its disabling effects can parallel those of deafness, particularly as approximately 90% of patients also have hearing loss⁵.

Despite its significant prevalence and impact, tinnitus lacks standardized treatment approaches, largely due to complex and multifactorial pathophysiology involving maladaptive neuroplasticity, central auditory system dysfunction, and oxidative stress^{1,3}. These mechanisms provide a foundation for exploring therapeutic interventions targeting oxidative stress, neuroinflammation, and neuronal hyperactivity¹. However, no drug has received approval from international regulatory bodies for tinnitus treatment^{4,5}. Current diagnosis relies on self-administered questionnaires, psychological evaluations, and assessments of perceptual characteristics^{3,6}. Management strategies primarily focus on symptom control through education, psychology-based approaches, sound therapy, hearing aids, and alternative and traditional therapies^{3,7}, while drug therapies primarily address comorbid symptoms like insomnia or anxiety^{3,8}.

In this context, many patients turn to alternative treatments, particularly plant extracts and dietary supplements, seeking relief from their symptoms^{9,10}. These interventions are chosen for their natural origin, accessibility, and lower cost compared to traditional treatments, though their safety and effectiveness are not always guaranteed^{3,9}. Plant extracts, whether derived from the entire plant or specific parts, are widely used; despite limited scientific evidence supporting their efficacy^{9,10}. Similarly, while approximately fifty-two dietary supplements have been reported for tinnitus treatment, including vitamin B₁₂, zinc, and melatonin⁹, scientific evidence supporting their efficacy remains limited^{9,10,11}, despite their popularity among older adults⁸.

While previous reviews have examined specific supplements like zinc and melatonin^{12,13}, no comprehensive systematic review has synthesized the evidence for both dietary

supplements and plant extracts in tinnitus treatment. This gap is particularly notable given the variety of interventions being studied, including Açai, traditional herbal medicines (e.g., Hangekobokuto), caffeine, and various dietary supplements (melatonin, alpha-lipoic acid, manganese, zinc, and vitamin B complex). Although previous research on vitamin B complex showed no superiority over placebo¹⁴, and a 2015 review suggested potential benefits of melatonin for tinnitus-related sleep disorders¹², the overall evidence landscape remains fragmented and unclear.

Therefore, this systematic review aims to comprehensively evaluate the efficacy of plant extracts and dietary supplements for tinnitus treatment, addressing a critical gap in current literature and providing evidence-based guidance for clinicians and patients.

METHODOLOGY

Study design and protocol registration

This systematic review and meta-analysis followed PRISMA guidelines¹⁵, with the protocol registered in PROSPERO (CRD42023446824).

Research question framework

The review question followed the PICO framework. The study population consisted of adults with acute or chronic tinnitus. Interventions encompassed herbal substances (e.g., Açai extract, caffeine) and dietary supplements (e.g., melatonin, alpha-lipoic acid, manganese, zinc sulfate, and vitamin B complex). Comparisons included placebo or concomitant interventions. Outcomes focused on tinnitus severity reduction, assessed through subjective measures including visual analogue scales (VAS), tinnitus loudness or pitch matching, and standardized self-reported questionnaires assessing the impact on health-related quality of life (HRQoL).

Search strategy

A systematic search was conducted in PubMed from inception through October 31, 2024. The search strategy, developed with a medical librarian, combined three conceptual blocks: tinnitus-related terms, dietary supplements and plant extracts interventions, and clinical trial methodology and outcomes assessment. Both Medical Subject Headings (MeSH) and free-text terms ([tiab]) were utilized, including specific terms for supplements (e.g., "melatonin", "zinc", "vitamins"), plant-based interventions (e.g., "herbal medicine", "plant extracts"),

and validated outcome measures (e.g., “Tinnitus Handicap Inventory”, “visual analog scale”). The complete search strategy is available in [Supplementary Table S1 \(https://www.renhyd.org/renhyd/article/view/2308/1305\)](https://www.renhyd.org/renhyd/article/view/2308/1305). Reference lists of eligible trials, systematic reviews, and meta-analyses were manually screened. The search was limited to English-language publications, with no date restrictions.

Inclusion criteria

Studies were included if they met four criteria: 1) design: randomized controlled trials (RCTs) or crossover trials (CO); 2) participants: adult patients with subjective tinnitus (either acute or chronic), or tinnitus due to sudden sensorineural hearing loss; 3) interventions: plant extracts (therapeutic plant-based substance or their derivatives, such as Açai, caffeine), or dietary supplements (including melatonin, alpha-lipoic acid, manganese, zinc, vitamin B complex); and 4) comparators: no intervention, placebo, or concomitant intervention control groups.

Outcome measures

The study included three outcome categories to evaluate the efficacy of plant extracts and dietary supplements on tinnitus: 1) subjective severity, measured through VAS for self-reported frequency or annoyance⁷; 2) psychoacoustic parameters, assessed through tinnitus loudness or pitch matching in decibels⁷; and 3) health-related quality of life (HRQoL), evaluated using standardized self-reported questionnaires including Tinnitus Handicap Inventory (THI), Tinnitus Functional Index (TFI), Tinnitus Handicap Questionnaire (THQ), or the Tinnitus Questionnaire (TQ)⁷. Detailed characteristics of assessment tools are provided in [Table S2 \(supplementary material, https://www.renhyd.org/renhyd/article/view/2308/1305\)](https://www.renhyd.org/renhyd/article/view/2308/1305).

Exclusion criteria

Studies were excluded if they met any of the following criteria: 1) non-RCT trials, or studies lacking a control group; 2) trial using Ginkgo biloba extracts, due to insufficient evidence for efficacy based on recent reviews, including an updated Cochrane review¹¹; 3) animal studies; 4) studies where tinnitus was a secondary outcome in populations with other primary conditions (e.g., dementia, Alzheimer’s disease); and 5) non-English publications.

Study selection process

The reviewers (YEC, GD) independently screened the titles and abstracts against predefined inclusion and exclusion criteria. Full texts of potentially eligible studies were then retrieved

and evaluated in detail. Disagreements were resolved through discussion, with a third author (RP) consulted when consensus could not be reached.

Data extraction

Two reviewers (YEC, LM) independently extracted data using standardized Excel forms. Extracted information included: first authors, publication year, recruitment country, study design, participant number, tinnitus type, treatment characteristics (dosage and administration methods), co-interventions, follow-up time, dropouts, adverse events, and outcome measurement. Disagreements were resolved through discussion.

Risk of bias assessment

Two reviewers (YE, RP) independently assessed risk of bias using the Revised Cochrane Risk-of-Bias Tool for Randomized Trials (RoB2)¹⁶. Assessment covered six domains: randomization process, deviations from intended interventions, missing outcome data, outcome measurement, selection of reported results, and other bias sources. Studies were classified as ‘low risk’ (all domains rated low), ‘some concerns’ (one or more domains with issues but none at high risk), or ‘high risk’ (significant methodological flaws). This classification system determined the overall quality of each study. Disagreements were resolved through consultation with a third author (LM).

Certainty of evidence assessment

We evaluated evidence certainty for each meta-analysis outcome using the modified GRADE (Grading, Development and Evaluation of Recommendations) approach. This framework assessed five domains that could reduce evidence certainty: risk of bias, inconsistency, indirectness, imprecision, and publication bias. Assessment method included: 1) inconsistency: I^2 statistic [$I^2 = ((Q - df)/Q) \times 100\%$], where Q is Cochran’s heterogeneity statistic and df is the degrees of freedom¹⁷; 2) publication bias: Egger’s test [$bias = \beta_0 + \beta_1 \times (1/SE)$], where SE is the standard error of the effect estimate¹⁸; 3) imprecision: optimal information size (OIS) calculation [$n = 2 \times (Z\alpha + Z\beta)^2 \times \sigma^2/\delta^2$], where $Z\alpha$ is the standard normal deviate for α , $Z\beta$ is the standard normal deviate for β , σ is the population standard deviation, and δ is the minimum clinically important difference^{19,20}. Evidence was ultimately rated as high, moderate, low, or very low certainty.

Statistical analysis

Analyses were conducted using RStudio (version 2022.07.02, RStudio PBC, Boston, MA, US) with the ‘metafor’ package. Random-effect models with restricted-maximum likelihood

(REML) estimation were employed for all analyses. For continuous outcomes, we calculated standardized mean differences (SMDs)¹⁸ with 95% confidence intervals (95%CI) using bias-corrected Hedges' g ¹⁸. For binary outcomes, odds ratios (OR) and their 95%CI were computed. Forest plots were generated to visualize treatment effects.

Statistical heterogeneity was assessed using I^2 statistics for between-study variance (I^2), τ^2 for absolute heterogeneity, and Q statistic (Q) for heterogeneity significance¹⁸. Subgroup analyses were conducted where possible. We performed sensitivity analyses using the leave-one-out method. Publication bias was evaluated through funnel plot inspection and the trim-and-fill method.

RESULTS

Study selection

The bibliographic search in PubMed identified 818 records. One additional record was identified through citation searching. After checking for duplicates, no duplicate records were found, and 818 articles underwent title and abstract screening. Following this process, 21 studies fulfilled all inclusion criteria and were selected for the final review (Figure 1 and [supplementary material, https://www.renhyd.org/renhyd/article/view/2308/1305](https://www.renhyd.org/renhyd/article/view/2308/1305)).

Study characteristics

The characteristics of the 21 articles are summarized in Table 1. The trials, published between 1987 and 2024, showed uneven geographic distribution: 28.6% (6 studies) from the USA (4)^{21–24} and Brazil (2)^{25,26}; 42.8% (9 studies) in Europe (including the United Kingdom (1)²⁷, Greece (3)^{28–30}, Spain (1)³¹, Denmark (1)³², the Netherlands (1)¹³, Italy (1)³³, and Romania (1)³⁴); 19.0% (4 studies) in the Middle East region (Turkey (1)³⁵ and Iran (3)^{36–38}); and 9.6% (2 studies) in Asia (India (1)³⁹, Japan (1)⁴⁰). No studies were identified from Africa or Oceania. All studies were RCTs, with four being CO (RCTs)^{20,21,23,26}. The trials included 1,759 participants (mean 83.8±50.3 per study; range 30 to 240), with mean age of 53.8 years (±9.9; IQR 7.7; range: 22–67.5). While gender distribution details are available in Table S3, comorbidity reporting was inconsistent across studies.

In tinnitus assessment, THI was the predominant tool for HRQoL measurement (47.6%, 10 studies), with additional use of THQ and TSI. For subjective perception, the VAS for loudness (19%, 4 studies) and for annoyance in one study. Psychoacoustic outcomes using pitch-match procedures in 14.3% (3) of trials.

Interventions fell into two categories: plant extracts and dietary supplements. Plant extracts were investigated in 5 trials: *Euterpe oleracea* (Açaí) extract²⁴, Hangekobokuto³⁶, Neurotec³⁵, and caffeine^{25,26}. Dietary supplements were examined in 16 studies, including: melatonin^{21,23,29,31,32,34}, zinc^{20,30,33}, alpha-lipoic acid²⁸, manganese with Lipoflavonoid Plus²², vitamin B (B_1 , B_6 , and B_{12})^{27,38,39}, and vitamin B₃ (nicotinamide)¹³, and MemoVigor Z³⁰. Treatment durations ranged from 0.1 to 24 weeks (median 8 weeks, IQR 8.0).

Clinical improvement parameters were defined in 50% of studies, typically as a 20-point THI reduction. No serious adverse events were reported ([see supplementary Table S4](#)).

Risk of bias assessment with RoB2

Overall, 57.1% (12/21) of studies showed 'some concerns' (7/12) or 'high risk' (5/12) of bias ([Figure S1](#)). The randomization process was the most affected domain due to poor reporting of randomization and allocation concealment, followed by missing outcome data due to high dropout rates. As shown in the summary graph, at the bottom of Figure S1, while some studies demonstrated low risk across multiple domains, the quality of methodological reporting varied considerably, with better control in outcome measurement and result selection domains.

FINDINGS FROM THE META-ANALYSIS

Effects of caffeine and melatonin on HRQoL and tinnitus loudness

For HRQoL, caffeine showed no significant effect compared to placebo in both common and random effects models (SMD=0.07; 95%CI: -0.73 to 0.86; $p=0.706$) with low heterogeneity ($I^2=0.0\%$; $Q=0.14$; $p=0.706$) across two studies^{25,26} ($n=73$ per group) (Figure 2-A). For melatonin interventions (Figure 2-B), three subgroup analyses were performed: combined therapy ($n=95$ intervention, $n=96$ control) showed a non-significant effect (MD=-2.43; 95%CI: -4.72 to -0.14; $I^2=64.1\%$); melatonin versus placebo ($n=57$ intervention, $n=57$ control) demonstrated no significant difference (MD=4.53; 95%CI: -1.85 to 10.90; $I^2=21\%$); and melatonin versus other drugs ($n=35$ intervention, $n=35$ control) showed no significant effect (MD=-6.67; 95%CI: -17.21 to 3.87)^{24,33,34,36}. For tinnitus loudness as a binary outcome (Figure 2-C), melatonin showed a significant reduction compared to placebo (OR of 2.52; 95%CI: 1.33–4.79; $p=0.005$), and low heterogeneity ($I^2=0.00$; $Q=0.10$; $p=0.755$) for the two trials^{21,23}.

Dropout rate and publication bias

The pooled dropout rate across 21 studies ($n=1,759$) was 19% (95%CI: 17%–22%) using a random effect model, with

Figure 1. Flowchart of the search and selection of papers on efficacy of plant extracts and dietary supplements for tinnitus (published before 31 October 2024) according to PRISMA guidelines.




Table 1. Characteristics of the included studies.

Author [reference] Year Study location	Design and setting	Sample size T:C (n)	Intervention protocol	Control/ comparisons	Treatment duration in weeks	Outcome's measurement	Dropout rate (%)	Risk of bias assessment
Abtahi <i>et al.</i> [34] 2017 Iran	RCT, parallel arms, DB Hospital center	(n=70) 35:35	Melatonin (3 mg/evening)	Sertraline (50 mg/evening)	12	THI*	4.29	Some concerns
Albu <i>et al.</i> [34] 2014 Romania	RCT, parallel arms, DB University hospital	(n=60) 30:30	Melatonin 3 mg + four intratympanic (IT) dexamethasone injections	Melatonin (3 mg) + IT placebo	12	THI*, (VAS) loudness, (VAS) awareness	8.33	Low
Arda <i>et al.</i> [33] 2003 Turkey	RCT, parallel arms, DB Hospital center	(n=41) 28:13	Zinc (50 mg/day)	Placebo	8	(VAS) loudness*, Tinnitus matching	0.00	Some concerns
Balatsouras <i>et al.</i> [30] 2024 Greece	RCT, Parallel arms, DB Hospital center	(n=240) 120:120	900 mg of MemoVigor 2 (Ginkgo biloba 3%, Bilberry 1%, phospholipids, vitamins B ₁ /B ₆ /B ₁₂ /C/E, selenium, magnesium, potassium, L-acetylcarnitine) once daily	Placebo	12	THI*	15.0	Low
Claire <i>et al.</i> [26] 2010 United Kingdom	RCT ^a , CO, DB University center studies	(n=66) 33:33	Days 1-15, caffeine- containing tea and/or coffee supplied (maintaining usual caffeine consumption). Days 16-18, phased withdrawal (1/2, 1/4 then 1/8 th normal caffeine consumption)	Days 19-30, decaffeinated supplies	4	TQ*, (VAS) annoyance, (VAS) frequency	1970	High
Coelho <i>et al.</i> [20] 2013 USA	RCT, CO, DB Tinnitus clinic	(n=116) 58:58	Zinc (50 mg/day) followed by a washout period of 30 days	Placebo	16	THQ*, (VAS) loudness, (VAS) annoyance	23.28	Low
Dadgarnia <i>et al.</i> [38] 2024 Iran	RCT, parallel arms, DB Hospital center	(n=140) 70:70	Vitamin B ₁₂ (500 microgram/day)	Placebo	4	THI*, VAS	0.00	Some concerns


Author [reference] Year Study location	Design and setting	Sample size T:C (n)	Intervention protocol	Control/ comparisons	Treatment duration in weeks	Outcome's measurement	Dropout rate (%)	Risk of bias assessment
Hulshof et al. [13] 1987 Netherlands	RCT, parallel arms, DB University hospital	(n=48) 24:24	Nicotinamide (B ₃ vitamin) capsules (70 mg/day)	Placebo	4	(VAS) annoyance*, Tinnitus matching	0.00	Some concerns
Hurtuk et al. [21] 2011 USA	RCT, CO, DB University center	(n=84) 42:42	Melatonin (3 mg/evening) washout period of 30 days	Placebo	4	TSI*, PSQL, BDI	27.38	Low
Ino et al. [36] 2013 Japan	RCT, parallel arms, DB University hospital	(n=76) 38:38	12 tablets twice a day (<i>Pinellia</i> tuber 6.0 g/day; <i>Poria sclerotium</i> , 5.0 g/day; <i>Magnolia</i> bark, 3.0 g/day; <i>Pirrellia</i> herb, 2.0 g/day; Zingiber rhizome 1.3 g/day)	Placebo	12	THI*, (VAS) loudness, (VAS) annoyance, SF-36	0.00	Some concerns
Khosravi et al. [35] 2023 Iran	RCT, parallel arms, DB Hospital center	(n=150) 80:70	100 mg of Neurotec (<i>Rosa canina</i> , <i>Urtica dioica</i> , <i>Tanacetum vulgare</i>) twice a day	Placebo	12	PTAs*, THI, (VAS) loudness, (VAS) annoyance	31.33	Low
Ledesma et al. [25] 2021 Brazil	RCT, parallel arms, TB University hospital	(n=80) 40:40	Caffeine (300 mg)	Placebo	Same day: evaluation one hour later	THI*, (VAS) loudness, Tinnitus matching	0.00	Low
Lopez-Gonzalez et al. [29] 2007 Spain	RCT, parallel arms, DB University hospital	(n=120) 30:30:30:30	Sulpiride (150 mg/day) or Melatonin (3 mg/evening) or sulpiride (150 mg/day) + melatonin (3 mg/evening)	Placebo	4	(VAS) loudness*, (VAS) perception: increase to disappearance	17.50	Some concerns
Markouet al. [27] 2004 Greece	RCT, parallel arms, DB University hospital	(n=108) 24:24:24:36	Trimetazidine (60 mg) + prednisolone (15 mg) + vitamin B complex (B ₁ , B ₆ , B ₁₂) or trimetazidine 60 mg or prednisolone + vitamin B complex (60 mg)	Piracetam (12 g within the first 30 minutes followed by another 12g during 24/h drip infusions + prednisolone (75 mg)	6	Tinnitus questionnaire modified*, acoustic gain	0.00	Some concerns
Neri et al. [31] 2009 Italy	RCT, parallel arms, DB University hospital	(n=102) 34:34:34	Sulodexide (250 mg/day) + Melatonin (3 mg/evening) or melatonin (3 mg/evening)	Placebo	12	THI*, Tinnitus matching	0.00	Some concerns

Author [reference] Year Study location	Design and setting	Sample size T:C (n)	Intervention protocol	Control/ comparisons	Treatment duration in weeks	Outcome's measurement	Dropout rate (%)	Risk of bias assessment
Oppitz <i>et al.</i> [24] 2022 Brazil	RCT, parallel arms, DB University hospital	(n=30) 15:15	Açaí extract (100 mg/day)	Placebo	12	THI*, BDI	0.00	Some concerns
Paaske <i>et al.</i> [30] 1991 Denmark	RCT, parallel arms, DB University hospital	(n=48) 23:25	Zinc (100 mg/day)	Placebo	8	(VAS) loudness*, Tinnitus matching	0.00	Some concerns
Petridou <i>et al.</i> [28] 2019 Greece	RCT, parallel arms, DB NDA	(n=70) 35:35	1 multivitamin/multimineral tablet/day + alpha-lipoic acid (600 mg/day)	Placebo	12	Tinnitus matching*, THI, TFI, (VAS) annoyance	10.00	Low
Rojas-Roncancio <i>et al.</i> [22] 2016 USA	RCT, parallel arms, DB University center	(n=40) 20:20	Manganese (8 mg/day) + Lipoflavonoid Plus pills three times a day	Lipoflavonoid Plus	24	Tinnitus matching*, THQ, (VAS) loudness	30.00	Some concerns
Rosenberg <i>et al.</i> [23] 1998 USA	RCT, CO, DB Hospital	(n=30) 15:15	Melatonin (3 mg/evening) washout period of 7 days	Placebo	4	THI*, tinnitus matching	23.33	Some concerns
Singh <i>et al.</i> [39] 2016 India	RCT, parallel arms, DB Medical College	(n=40) 20:20	Vitamin B ₁₂ (2500 mcg/week IM)	Placebo (isotonic saline 1ml IM)	6	THI*, VAS	0	Some concerns

BDI: Beck depression inventory; **CO:** Crossover trial; **DB:** Double-blind; **mcg:** Micrograms; **n:** Number of participants for each group; **NDA:** No data available;
PSQI: Pittsburgh sleep quality index; **PTAs:** Pure-tone audiograms; **RCT:** Randomized controlled trial; **SF-36:** Short Form Health Survey 36;
T:C (n): Number of participants in treatment and control groups; **TB:** Triple-blind; **THI:** Tinnitus Handicap Inventory; **THQ:** Tinnitus Handicap Questionnaire;
Q: Tinnitus Questionnaire; **TSI:** Tinnitus Severity Index; **VAS:** Visual analogue scale.

Note: [a] Pseudo-randomized trial; [b] Primary outcome.

Figure 2. Metanalysis of the effect of caffeine on HRQoL (A), melatonin on HRQoL (B), and melatonin on tinnitus loudness (C). Forest plot showing odd ratios (OR) and 95%CI. The risk of bias was assessed using RoB2 tool across six domains using traffic light system.



substantial heterogeneity ($I^2=79.9\%$; $\tau^2=1.9662$; $p<0.0001$). Individual rates ranged from 0% to 31% (Khosravi et al.) and 30% (Rojas-Roncancio et al.), as detailed in Figure S2. Funnel plots

(Figure S3) showed asymmetric distribution patterns across all comparisons, although trim-and-fill analysis did not impute any studies for caffeine and melatonin effects. Formal publication

bias assessment via Egger’s test was not feasible due to limited studies per comparison (<10). Leave-one-out sensitivity analysis (Figure S4) confirmed stability of effect estimates when removing individual studies.

Certainty of evidence

All outcomes were rated as low-certainty evidence (●●○○) using the GRADE system (Table 2). For HRQoL, neither caffeine (SMD: 0.07, 95%CI: -0.73 to 0.86) nor melatonin showed significant effects compared to placebo. Melatonin demonstrated a significant improvement on tinnitus loudness (OR: 2.52, 95%CI: 1.33 to 4.79). Evidence was downgraded due to serious risk of bias (as shown in the risk of bias panels of Figures 2-4) and imprecision from small sample size.

DISCUSSION

This systematic review and meta-analysis evaluated the efficacy of plant extracts and dietary supplements on tinnitus, analyzing 21 studies (1,759 adults) across three key outcomes: subjective perception, HRQoL, and perceived loudness. The GRADE assessment rated all meta-analyzed outcomes as low-certainty evidence (●●○○), primarily due to serious risk of bias and imprecision from small sample sizes. Despite low statistical heterogeneity ($I^2=0\%$), the limited number of trials (<3 per intervention) and small sample sizes prevented robust

assessment of publication bias and reduced confidence in effect estimates. Results varied across studies due to diverse outcome measures and differences in participants’ chronic tinnitus conditions. The evidence synthesis focused on nine interventions: four plant extracts and five dietary supplements, with only melatonin and caffeine providing sufficient data for meta-analysis.

Meta-analysis of two crossover trials (164 participants) demonstrated that melatonin modestly reduces tinnitus loudness (OR: 2.52, 95%CI: 1.33 to 4.79)^{21,23}. Both trials used similar protocols (30-day treatment, 3 mg melatonin dosage), with adequate washout periods given melatonin’s short half-life (10-60 minutes post-oral administration)⁴¹. Although melatonin may improve sleep quality in chronic subjective tinnitus patients^{36,42}, its long-term clinical remains uncertain. Regarding caffeine, our quantitative synthesis showed no efficacy in reducing the impact of tinnitus on HRQoL (SMD: 0.07, 95%CI: -0.26 to 0.39). This aligns with previous reviews⁴³, showing mixed results, where some individuals reported temporary relief^{8,44}, but larger studies failed to establish a clear benefit⁸. The relationship between caffeine and tinnitus remains complex, warranting more rigorous investigation.

Additionally, evidence for other interventions remains limited. Single studies of plant extracts (Açaí extract²⁴, Hangekobokuto compound³⁶, and Neurotec compound³⁵, alpha-lipoic acid²⁸, manganese²² and MemoVigor 2³⁰) preclude definitive conclusions about their efficacy. Furthermore, trials using a vitamin B complex in combination with drugs²⁷ showed no

Table 2. GRADE summary of findings.

Outcome	Intervention	Comparator	Pooled effect (95%CI)	Studies (n)	Study design	Certainty of evidence	Downgrade reason
HRQoL	Caffeine	Placebo	SMD: 0.07 (-0.73; 0.86)	2 (146)	RCT	●●○○ Low	• Risk of bias ¹ • Imprecision ²
HRQoL	Melatonin	Placebo	MD: 4.53 (-1.85; 10.90)	2 (114)	RCT	●●○○ Low	• Risk of bias ¹ • Imprecision ²
Tinnitus Loudness	Melatonin	Placebo	OR: 2.52 (1.33; 4.80)	2 (168)	RCT	●●○○ Low	• Risk of bias ¹ • Imprecision ²

Certainty of Evidence Legend: ●●●● High; ●●●○ Moderate; ●●○○ Low; ●○○○ Very Low.

HRQoL: Health-related quality of life; SMD: Standardized mean difference; MD: Mean difference; OR: Odds ratio; n: Number of participants; RCT: Randomized controlled trial; 95%CI: 95% confidence interval.

Footnotes: ¹ Downgraded one level due to serious risk of bias (multiple domains showing some concerns or high risk); ² Downgraded one levels due to imprecision (number of studies < 3; Wide confidence intervals).

evidence of effect on subjective discomfort or HRQoL. Studies of B₁₂^{38,39} and vitamin B₃¹³ similarly demonstrated no significant benefits, although methodological heterogeneities prevented meta-analysis.

These findings align with current evidence and clinical guidelines. While various supplements have been explored for tinnitus management, including zinc, manganese, and B vitamins, results remain inconsistent^{3,44}. Recent recommendations^{8,45} and the European multidisciplinary guideline for tinnitus management specifically caution against using dietary supplements and herbal medicines as primary treatments, citing lack of proven efficacy and potential risks³. Although certain supplements may benefit specific conditions (e.g., zinc deficiency, sleep disturbances), they should not be recommended as standalone treatments^{44,46}.

Methodological heterogeneity across studies posed significant challenges for evidence synthesis. Outcome measures varied considerably, with two Visual Analog Scales (VAS-loudness and VAS-annoyance) assessing subjective tinnitus perception and five questionnaires evaluating HRQoL impact (THI, TSI, THQ, TQ, TQ-SF), each with distinct sensitivities to tinnitus domains^{6,47}. While tinnitus matching could provide sensitive endpoint measures⁴, only 14% of studies reported these outcomes. Additionally, despite focusing on subjective chronic tinnitus⁷, only six established minimum tinnitus scores as inclusion criteria. The geographical concentration of trials in Europe and North America, with limited representation from Asia, Africa, and Oceania, further constrains the generalizability of findings, as populations from underrepresented regions might differ in terms of healthcare access, dietary patterns, and genetic factors affecting treatment responses^{2,4}.

Future directions, strengths and limitations

Future studies should adhere to CONSORT guidelines, particularly regarding randomization, allocation concealment, and blinding. Studies should establish minimum threshold for tinnitus loudness using validated measurement tools. The Unification of Treatments and Interventions for Tinnitus Patients Trial⁴⁸ offers standardized methodologies that could enhance study comparability, emphasizing longer follow-up periods and the consideration of common comorbidities like insomnia, anxiety, and depression.

Understanding tinnitus mechanisms remains crucial, given limited evidence on tinnitus risk factors⁴⁹. Novel approaches, such as machine learning and random forest regression⁵⁰, show promise for predicting optimal interventions but requires validation in larger, diverse populations. Recent findings on problematic tinnitus predictors⁵¹ emphasize the importance of identifying subgroups for tailored interventions.

Despite our comprehensive search strategy, this review has several limitations. The exclusion of non-English publications, varying methodological quality across studies, and geographical concentration in Europe and North America limit global applicability. Inconsistent reporting of comorbidities and the limited number of studies prevented us from performing robust subgroup analyses and publication bias assessment, though the trim-and-fill method and funnel plots were performed.

CONCLUSIONS

The systematic review and meta-analysis found low-certainty evidence that melatonin may slightly reduce tinnitus loudness, while caffeine showed no effect on tinnitus symptoms or HRQoL. Evidence for other plants extracts and dietary supplements remains insufficient due to limited studies and methodological limitations. The GRADE assessment revealed serious risk of bias and imprecision across studies, primarily due to small sample sizes. These findings support current clinical practice guidelines that advise against dietary supplements and herbal medicines as primary treatments for tinnitus. While specific supplements might offer targeted benefits (e.g., melatonin for sleep disturbances), their routine use in tinnitus management cannot be recommended based on current evidence.

AUTHORS' CONTRIBUTIONS

Y.E.C., G.D.G., L.M.A., and R.P.M. contributed to the study's conceptualization, design, data collection, analysis, and interpretation. All authors were involved in the methodology and initial article drafting. R.P.M., L.M.A. and G.D.G. revised the draft. Y.E.C. completed the final editing. All authors reviewed and approved the final version of the manuscript.

FUNDING

This study received no external funding.

DATA AVAILABILITY STATEMENT

Data supporting this study's findings are available from the corresponding author upon reasonable request.

COMPETING INTERESTS

The authors declare that there were no conflicts of interest.

REFERENCES

- (1) Langguth B, De Ridder D, Schlee W, Kleinjung T. Tinnitus: Clinical Insights in Its Pathophysiology-A Perspective. *J Assoc Res Otolaryngol*. 2024; 25(3): 249-58. doi: 10.1007/s10162-024-00939-0.
- (2) Jarach CM, Lugo A, Scala M, van den Brandt PA, Cederroth CR, Odone A, et al. Global Prevalence and Incidence of Tinnitus: A Systematic Review and Meta-analysis. *JAMA Neurol*. 2022; 79(9): 888-900. doi: 10.1001/jamaneurol.2022.2189.
- (3) Cima RFF, Mazurek B, Haider H, Kikidis D, Lapira A, Noreña A, et al. A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment. *Hno*. 2019; 67(1): 10-42.
- (4) Cederroth CR, Gallus S, Hall DA, Kleinjung T, Langguth B, Maruotti A, et al. Editorial: Towards an Understanding of Tinnitus Heterogeneity. *Front Aging Neurosci*. 2019; 11.
- (5) Bhatt JM, Lin HW, Bhattacharyya N. Prevalence, Severity, Exposures, and Treatment Patterns of Tinnitus in the United States. *JAMA Otolaryngol Neck Surg*. 2016; 142(10): 959-65. doi: 10.1001/jamaoto.2016.1700.
- (6) Hall DA, Haider H, Szczepek AJ, Lau P, Rabau S, Jones-Diette J, et al. Systematic review of outcome domains and instruments used in clinical trials of tinnitus treatments in adults. *Trials*. 2016; 17(1): 270. doi: 10.1186/s13063-016-1399-9.
- (7) Okayasu T, Mitani K, Kitahara T. Reviewing Kampo medicine (Traditional Japanese Herbal Medicine) for otology/neurotology diseases. *Auris Nasus Larynx*. 2024; 51(1): 25-30. doi: 10.1016/j.anl.2023.04.009.
- (8) Lee HY, Jung DJ. Recent Updates on Tinnitus Management. *J Audiol Otol*. 2023; 27(4): 181-92. doi: 10.7874/jao.2023.00416.
- (9) Coelho C, Tyler R, Ji H, Rojas-Roncancio E, Witt S, Tao P, et al. Survey on the Effectiveness of Dietary Supplements to Treat Tinnitus. *Am J Audiol*. 2016; 25(3): 184-205. doi: 10.1044/2016_AJA-16-0021.
- (10) Alonso Street EM, Grutzmacher S, Branscum A, Smit E. Self-Perceived Health and Chronic Conditions Among Users of Supplements With and Without Botanical Ingredients: Findings from the 2009–2014 National Health and Nutrition Examination Survey. *J Diet Suppl*. 2022; 19(6): 689-703.
- (11) Sereda M, Xia J, Scutt P, Hilton MP, El Refaie A, Hoare DJ. Ginkgo biloba for tinnitus. *Cochrane Database Syst Rev*. 2022; 2023(1). doi: 10.1002/14651858.CD013514.pub2.
- (12) Miroddi M, Bruno R, Galletti F, Calapai F, Navarra M, Gangemi S, et al. Clinical pharmacology of melatonin in the treatment of tinnitus: a review. *Eur J Clin Pharmacol*. 2015; 71(3): 263-70. doi: 10.1007/s00228-015-1805-3.
- (13) Person OC, Puga ME, da Silva EM, Torloni MR. Zinc supplementation for tinnitus. *Cochrane Database Syst Rev*. 2016; 2016(11). doi: 10.1002/14651858.CD009832.pub2.
- (14) Hulshof JH, Vermeij P. The effect of nicotinamide on tinnitus: a double-blind controlled study. *Clin Otolaryngol*. 1987; 12(3): 211-4. doi: 10.1111/j.1365-2273.1987.tb00189.x.
- (15) Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*. 2015; 349(jan02 1): g7647-g7647. doi: 10.1136/bmj.g7647.
- (16) Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019; 14898. doi: 10.1136/bmj.14898.
- (17) Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003; 327(7414): 557-60.
- (18) Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997; 315(7109): 629-34. doi: 10.1136/bmj.315.7109.629.
- (19) Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence—imprecision. *J Clin Epidemiol*. 2011; 64(12): 1283-93.
- (20) Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane Handbook for Systematic Reviews of Interventions*. 2nd ed. Newark: John Wiley & Sons, Incorporated; 2019.
- (21) Coelho C, Witt SA, Ji H, Hansen MR, Gantz B, Tyler R. Zinc to Treat Tinnitus in the Elderly: A Randomized Placebo Controlled Crossover Trial. 2013; 34(6).
- (22) Hurtuk A, Dome C, Holloman CH, Wolfe K, Welling DB, Dodson EE, et al. Melatonin: Can it Stop the Ringing? *Ann Otol Rhinol Laryngol*. 2011; 120(7): 433-40. doi: 10.1177/000348941112000703.
- (23) Rojas-Roncancio E, Tyler R, Jun H-J, Wang T-C, Ji H, Coelho C, et al. Manganese and Lipoflavonoid Plus® to Treat Tinnitus: A Randomized Controlled Trial. *J Am Acad Audiol*. 2016; 27(08): 661-8. doi: 10.3766/jaaa.15106.
- (24) Rosenberg SI, Silverstein H, Rowan PT, Olds MJ. Effect of Melatonin on Tinnitus. *The Laryngoscope*. 1998; 108(3): 305-10. doi: 10.1097/00005537-199803000-00001.
- (25) Oppitz SJ, Garcia MV, Bruno RS, Zemolin CM, Baptista BO, Turra BO, et al. Supplementation with açai (Euterpe Oleracea Martius) for the treatment of chronic tinnitus: effects on perception, anxiety levels and oxidative metabolism biomarkers. *CoDAS*. 2022; 34(4): e20210076. doi: 10.1590/2317-1782/20212021076.
- (26) Ledesma ALL, Leite Rodrigues D, Monteiro De Castro Silva I, Oliveira CA, Bahmad F. The effect of caffeine on tinnitus: Randomized triple-blind placebo-controlled clinical trial. *PLOS ONE*. 2021; 16(9): e0256275. doi: 10.1371/journal.pone.0256275.
- (27) Claire LSt, Stothart G, McKenna L, Rogers PJ. Caffeine abstinence: an ineffective and potentially distressing tinnitus therapy. *Int J Audiol*. 2010; 49(1): 24-9. doi: 10.3109/14992020903160884.
- (28) Markou K, Nikolaou A, Petridis DG, Vlachtsis KC, Kouloulas A, Daniilidis IC. Evaluation of various therapeutic schemes in the treatment of tinnitus due to acute acoustic trauma. *Kulak Burun Bogaz Ihtis Derg KBB J Ear Nose Throat*. 2004; 12(5-6): 107-14.
- (29) Petridou AI, Zagora ET, Petridis P, Korres GS, Gazouli M, Xenelis I, et al. The Effect of Antioxidant Supplementation in Patients with Tinnitus and Normal Hearing or Hearing Loss: A Randomized, Double-Blind, Placebo Controlled Trial. *Nutrients*. 2019; 11(12): 3037. doi: 10.3390/nu11123037.
- (30) Balatsouras DG, Papitsi I, Koukoutsis G, Katotomichelakis M. The effect of MemoVigor 2 on recent-onset idiopathic tinnitus: a randomized double-blind placebo-controlled clinical trial. *Front Pharmacol*. 2024; 15: 1252343. doi: 10.3389/fphar.2024.1252343.
- (31) Lopez-Gonzalez MA, Santiago AM, Esteban-Ortega F. Sulpiride

- and Melatonin Decrease Tinnitus Perception Modulating the Auditolimbic Dopaminergic Pathway. *J Otolaryngol.* 2007; 36(04): 213. doi: 10.2310/7070.2007.0018.
- (32) Paaske PB, Kjems G, Pedersen CB, Sam ILK. Zinc in the Management of Tinnitus: Placebo-Controlled Trial. *Ann Otol Rhinol Laryngol.* 1991; 100(8): 647-9. doi: 10.1177/000348949110000809.
- (33) Neri G, De Stefano A, Baffa C, Kulamarva G, Di Giovanni P, Petrucci G, et al. Treatment of central and sensorineural tinnitus with orally administered Melatonin and Sulodexide: personal experience from a randomized controlled study. *Acta Otorhinolaryngol Ital Organo Uff Della Soc Ital Otorinolaringol E Chir Cerv-Facc.* 2009; 29(2): 86-91.
- (34) Albu S, Chirtes F. Intratympanic dexamethasone plus melatonin versus melatonin only in the treatment of unilateral acute idiopathic tinnitus. *Am J Otolaryngol.* 2014; 35(5): 617-22. doi: 10.1016/j.amjoto.2014.06.009.
- (35) Arda HN, Tuncel U, Akdogan O, Ozluoglu LN. The Role of Zinc in the Treatment of Tinnitus: *Otol Neurotol.* 2003; 24(1): 86-9. doi: 10.1097/00129492-200301000-00018.
- (36) Abtahi S, Hashemi S, Mahmoodi M, Nilfroush M. Comparison of melatonin and sertraline therapies on tinnitus: A randomized clinical trial. *Int J Prev Med.* 2017; 8(1): 61. doi: 10.4103/ijpvm.ijpvm_229_17.
- (37) Khosravi MH, Atefi A, Mehri A, Sodeifian F, Yousefi J, Bagheri Hagh A, et al. Therapeutic effects of *Rosa canina*, *Urtica dioica* and *Tanacetum vulgare* herbal combination in treatment of tinnitus symptoms: A double-blind randomised clinical trial. *Clin Otolaryngol.* 2023; 48(2): 151-7. doi: 10.1111/coa.13989.
- (38) Dadgarnia M, Mandegari M, Zand V, Vaziribozorg S, Zare-Zardini H, Mousaviyazdi K, et al. The effect of vitamin B12 on idiopathic tinnitus. *Am J Otolaryngol.* 2024; 45(1): 104028. doi: 10.1016/j.amjoto.2023.104028.
- (39) Singh C, Kawatra R, Gupta J, Awasthi V, Dungana H. Therapeutic role of Vitamin B12 in patients of chronic tinnitus: A pilot study. *Noise Health.* 2016; 18(81): 93-7. doi: 10.4103/1463-1741.178485.
- (40) Ino T, Odaguchi H, Wakasugi A, Oikawa T, Sano H, Okamoto M, et al. A randomized, double-blind, placebo-controlled clinical trial to evaluate the efficacy of hangekobokuto in adult patients with chronic tinnitus. *J Tradit Med.* 2013; 30(2): 72-81. doi: 10.11339/jtm.30.72.
- (41) Minich DM, Henning M, Darley C, Fahoum M, Schuler CB, Frame J. Is Melatonin the "Next Vitamin D"? A Review of Emerging Science, Clinical Uses, Safety, and Dietary Supplements. *Nutrients.* 2022; 14(19): 3934. doi: 10.3390/nu14193934.
- (42) Chen J-J, Chen Y-W, Zeng B-Y, Hung C-M, Zeng B-S, Stubbs B, et al. Efficacy of pharmacologic treatment in tinnitus patients without specific or treatable origin: A network meta-analysis of randomised controlled trials. *EClinicalMedicine.* 2021; 39: 101080. doi: 10.1016/j.eclinm.2021.101080.
- (43) Hofmeister M. Do dietary factors significantly influence tinnitus? *Aust J Gen Pract.* 2019; 48(3): 153-7. doi: 10.31128/AJGP-07-18-4643.
- (44) DiSogra RM, O'Brien-Russo CA, Deshpande AK. Dietary Supplements, Essential Oils, and Cannabinoids for Tinnitus Relief. *Tinnitus: Advances in Prevention, Assessment, and Management.* San Diego, CA: Plural Publishing; 2022. p. 247-60.
- (45) Meijers S, Stegeman I, Van Der Leun JA, Assegaf SA, Smit AL. Analysis and comparison of clinical practice guidelines regarding treatment recommendations for chronic tinnitus in adults: a systematic review. *BMJ Open.* 2023; 13(9): e072754. doi: 10.1136/bmjopen-2023-072754.
- (46) Ashan SF, Tyler RS. Use of Supplements for Management of Botherome Subjective Tinnitus. 2019.
- (47) Genitsaridi E, Hoare DJ, Kypraios T, Hall DA. A Review and a Framework of Variables for Defining and Characterizing Tinnitus Subphenotypes. *Brain Sci.* 2020; 10(12): 938. doi: 10.3390/brainsci10120938.
- (48) Simoes JP, Schoisswohl S, Schlee W, Basso L, Bernal-Robledano A, Boecking B, et al. The statistical analysis plan for the unification of treatments and interventions for tinnitus patients randomized clinical trial (UNITI-RCT). *Trials.* 2023; 24(1): 472. doi: 10.1186/s13063-023-07303-2.
- (49) Biswas R, Genitsaridi E, Trpchevska N, Lugo A, Schlee W, Cederroth CR, et al. Low Evidence for Tinnitus Risk Factors: A Systematic Review and Meta-analysis. *J Assoc Res Otolaryngol.* 2022; 24(1): 81-94. doi: 10.1007/s10162-022-00874-y.
- (50) Bromis K, Sarafidis M, Manta O, Kouris I, Vellidou E, Schlee W, et al. Predicting the optimal therapeutic intervention for tinnitus patients using random forest regression: A preliminary study of UNITI's decision support system model. 2022 44th Annual International Conference of the IEEE Engineering in Medicine & Biology Society (EMBC). IEEE; 2022. p. 2655-8.
- (51) De Gruy JA, Lorenzo WW, Vu T-H, Paul O, Lee C, Spankovich C. Prevalence and predictors of problematic tinnitus. *Int J Audiol.* 2024; 1-7. doi: 10.1080/14992027.2024.2378804.