



Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics

INVESTIGACIÓN – **versión post-print**

Esta es la versión aceptada para publicación. El artículo puede recibir modificaciones de estilo y de formato.

Development and evaluation of Diet Quality Index for Brazilians based on the updated version of the 2024 Food Pyramid: Nutrition focus House Budget Survey, 2017-18

Desarrollo y evaluación del índice de calidad de la dieta para brasileños basado en la versión actualizada de la Pirámide Alimenticia de 2024: Encuesta de presupuesto interno con enfoque nutricional, 2017-18

Ana Carolina Barco Leme^{a,b,*}, Sonia Tucunduva Philippi^b

^a Faculty of Health Sciences, Universidad Autónoma de Chile, Metropolitan area, 7500912, Chile

^b Department of Nutrition, School of Public Health, University of São Paulo, São Paulo, 01246-904, Brazil.

*acarol.leme@gmail.com

Recibido: 17/01/2025; Aceptado: 04/04/2025; Publicado: 06/05/2025.

Editor Asignado: Tania Fernández Villa, Universidad de León, León, España

CITA: Barco Leme AC, Tucunduva Philippi S. Development and evaluation of Diet Quality Index for Brazilians based on the updated version of the 2024 Food Pyramid: Nutrition focus House Budget Survey, 2017-18. Rev Esp Nutr Hum Diet. 2025; 29(2). doi: 10.14306/renhyd.29.2.2393 [ahead of print].

La Revista Española de Nutrición Humana y Dietética se esfuerza por mantener a un sistema de publicación continua, de modo que los artículos se publiquen antes de su formato final (antes de que el número al que pertenecen se haya cerrado y/o publicado). De este modo, intentamos poner los artículos a disposición de los lectores/usuarios lo antes posible.

The Spanish Journal of Human Nutrition and Dietetics strives to maintain a continuous publication system, so that the articles are published before its final format (before the number to which they belong is closed and/or published). In this way, we try to put the articles available to readers/users as soon as possible.

ABSTRACT

Introduction: In 2024 the Brazilian Food Pyramid was updated based on 7 food groups that meet the current dietary recommendations and appeared to be useful to help population achieve healthy eating behaviours. As many epidemiologic studies use dietary indexes, the current study developed and evaluated the psychometric properties of a dietary index to measure adherence to the updated food pyramid groups recommendations on healthy food choices.

Methods: Dietary intake data from the mean of two 24-hour dietary recalls in the Nutrition Focus House Budget Survey, 2017-18 was used for this purpose. Variability among individuals were assessed via distribution of dietary intakes. Mean dietary scores were compared among population subgroups. Pearson's correlations were used to verify the relationship between each component, overall scores and total energy intake. Finally, multidimensionality was checked with principal components analysis based on the 9 components of the index.

Results: Mean Brazilian index score was 65.65/100 (95% CI, 65.55, 65.80) among Brazilians $\geq$ 10yo. Adolescents (64.36) and males' adults (63.94) had the lowest and female adults (66.84) had the highest total score. Sodium showed the strongest negative correlation with fruits and vegetables ($r = -0.52$), while total score the strongest positive correlation with fatty acids ($r = 0.52$). Energy intake showed a stronger positive correlation for grains ($r = 0.71$) and a negative for fruits and vegetables ($r = -0.21$). Principal components analysis revealed at least 4 dimensions (60% total variance).

Conclusions: The Brazilian Healthy Eating Index, 2024 showed good psychometric properties supporting the 2024 updated food pyramid recommendation.

Keywords: Dietary Guidelines; Diet, food and nutrition; Cross-sectional studies.

RESUMEN

Introducción: En 2024, la Pirámide Alimenticia Brasileña fue actualizada basándose en 7 grupos de alimentos que cumplen con las recomendaciones dietéticas actuales y parecen ser útiles para ayudar a la población a lograr conductas alimentarias saludables. Como muchos estudios epidemiológicos utilizan índices dietéticos, el estudio actual desarrolló y evaluó las propiedades

psicométricas de un índice dietético para medir la adherencia a las recomendaciones actualizadas de los grupos de la pirámide alimenticia sobre la elección de alimentos saludables.

Metodología: Para este propósito se utilizaron datos de ingesta dietética de la media de dos recordatorios dietéticos de 24 horas en Encuesta sobre el presupuesto de la Cámara de Enfoque en Nutrición, 2017-18. La variabilidad entre los individuos se evaluó mediante la distribución de la ingesta dietética. Se compararon las puntuaciones dietéticas medias entre subgrupos de población. Se utilizaron correlaciones de Pearson para verificar la relación entre cada componente, las puntuaciones generales y la ingesta total de energía. Finalmente, se comprobó la multidimensionalidad con un análisis de componentes principales basado en los 9 componentes del índice.

Resultados: La puntuación media del índice brasileño fue 65,65/100 (IC del 95 %, 65,55, 65,80) entre los brasileños ≥ 10 años. Los adolescentes (64,36) y los hombres adultos (63,94) tuvieron la puntuación total más baja y las mujeres adultas (66,84) la puntuación total más alta. El sodio mostró la correlación negativa más fuerte con las frutas y verduras ($r=-0,52$), mientras que la puntuación total la correlación positiva más fuerte con los ácidos grasos ($r=0,52$). La ingesta de energía mostró una correlación positiva más fuerte para los cereales ($r=0,71$) y negativa para las frutas y verduras ($r=-0,21$). El análisis de componentes principales reveló al menos 4 dimensiones (60% de varianza total).

Conclusión: El Índice Brasileño de Alimentación Saludable de 2024 mostró buenas propiedades psicométricas que respaldan la recomendación de la pirámide alimentaria actualizada de 2024.

Palabras clave: Pautas dietéticas, Dieta, alimentación y nutrición, estudios transversales.

KEY MESSAGES

- The food pyramid reflects the dietary recommendations of a target population.
- The psychometric properties of diet quality indices should be assessed.
- Variability and correlations between dietary components reflect good psychometrics.
- It is recommended to configure the weights of different components when targeting specific groups.

INTRODUCTION

There is clear evidence demonstrating the relationship between suboptimal dietary patterns and leading risk factors for morbidity and mortality in Brazil and worldwide. Therefore, healthy eating is important to promote health from earlier to later in life. Healthy eating is a dietary pattern favouring all type of food choices, not stigmatizing into “good” or “bad, promoting the complete physical, social and emotional well-being¹. Many epidemiological studies have examine the associations between health outcomes and single foods or nutrients, but they are consumed in combinations which can induce interactions and synergies between dietary components². Dietary pattern analysis is known to be a more appropriate approach in investigating health-related outcomes rather than focusing on single foods or nutrient³. Developing *a priori* dietary index is one approach to assess dietary patterns, i.e., pre-defined algorithm to quantify food and nutrient intake relative to national and international nutritional recommendations⁴. Indices can be used to measure diet quality in populations, monitoring over time or dietary changes in trials. Furthermore, in epidemiological studies an index can be used to investigate the associations with health and diet. Also, confounder factors can be controlled by using diet index score⁵. Several indexes have been developed, and a well-known example is the “American” Healthy Eating Index (HEI), which has passed through several updates to align to the US dietary guidelines². This index has been associated with health outcomes and as monitoring tool in North America populations⁵. More recently, the Global Diet Quality Score (GDQS)⁶ have been developed known as an “easy-to-be-used” tool and with an 25-item metric for nutrient adequacy and diet related diseases, more specifically for chronic non-communicable diseases. The GDQS have been correlated with the intake of “ultra-processed” foods and adequacy of some nutrients with the Brazilian population⁷. However, these indexes have limitations to be used in the Brazilian context. First, the “American” HEI is based on the US dietary guidelines and the diet is culturally different from Brazil. Second, the GDQS is a metric to focus on monitoring chronic non-communicable diseases not comprehending the overall concept of health. For instance, nutrition and diet-related health are driven by multiple factors and are embedded in social determinants of health and structural factors. These include poverty, lack of access to quality education and employment, poor quality housing, unfavourable work and neighborhood

conditions, transportation issues, environmental racism, neighborhood violence, and the clustering of disadvantaged groups^{8,9}.

In 2024, Philippi et al.¹⁰ updated an iconographic with seven food groups targeting overall health of the Brazilian population. Furthermore, this iconographic focused on the population health disparities that has been associated with factors at an individual level⁸, e.g., food (in) security, which is the (lack) consistent, dependable access to sufficient food for an active and healthy life¹¹. This factor can influence eating habits, and dietary preference and are themselves influenced by social and structural factors that contribute to diet-related disparities⁸. Therefore, addressing multiple intersecting factors that influence dietary patterns and their contribution to diet related health disparities is essential for developing effective tools to monitoring diet and health disparities.

To date, two Brazilian indices have been developed, both based on an iconographic that was endorsed by the 2006 Brazilian Food Guide¹². The first dietary quality index consisted of five criteria: < 30% of energy from total fats, < 10% of energy from saturated fats, < 300mg/day of cholesterol, $\leq$ 2400mg/day of sodium, and $\geq$ 5 servings of fruits and vegetables. For each criterion, scores ranged from 0 to 10 depending on adherence to recommendation. The score of (total and saturated) fats, cholesterol, and sodium was inversely related to the total diet score¹². The second index was an updated version of the previous one and consisted of 12 components. The score, ranging from zero (low quality) to 96 (high quality) was based on the (American) HEI 2005 version cut-off values.

Psychometric properties of an index measuring adherence to given set of dietary recommendations should be verified before the use in epidemiological, intervention and surveillance studies. Certainly, assessing adherence to recommendations in an iconographic based on food groups (also known as the Food Pyramid) with a validated index may reduce the risk for inconsistent research findings that may impact policy decision-making and lead to confusing messages to the public. Several dimensions of validity should be applicable to validate the index and include: (i) construct validity, i.e., extent to which a given index varies according to other variables known to be associated to diet quality; and (ii) reliability, i.e., extent to which all components of an index are internally consistent¹³. The aim of this study was to develop and

evaluate the construct validity and reliability in terms of internal consistency of a metric based on the updated Food Pyramid groups.

METHODS

Study design and participants

Data from the 2017-18 Nutrition focus House Budget Survey was used to conduct the analyses. The House Budget Survey is a nationally representative survey of individuals ≥ 10 yo living in private dwelling from the 26 Brazilian states plus the Federal District. Individuals from the military forces, penitentiaries, shelters, orphanages, hospitals and other institutionalized venues were excluded. Data were collected between August 11th of 2017 to August 11th of 2018. Pregnant and lactating women were excluded (less than 4% of all participants in the Nutrition focus House Budget Survey (n=707)). Analyses were based on the public use microdata files obtained from the Brazilian Geography and Statistics Institute. Secondary analyses of data from the Nutrition-focus House Budget Survey do not require additional ethics approval.

Dietary assessment

The 24h-recall was the method of choice to assess dietary intake of the population and followed the Automated Multiple Pass Method. The average of two non-consecutive dietary recalls were taken to estimate each individual dietary intake¹³. From the total sample of the Brazilian – House Budget Survey, 2017-18 (n = 178,431), 25% of the respondents completed the 1st 24h-recall (n = 46,164) and 22% the second (n = 38,854) via computer-assisted interviews by a trained personal. The Global Diet software was used to translate all the information related to foods, beverages, food preparations, amounts, methods of preparations, added items and meal occasions and venues. Reported dietary intakes measured via 24-hour recall were 1st classified according to the food categories used to calculate the group components of the dietary metric and were then summed per respondent for each 24h recall available. The nutrients used to calculate the metric, i.e., saturated, mono-unsaturated (MONO) and poli-unsaturated (POLI) fats, free sugar, and sodium, expressed in grams were deemed as reference amounts (RA). Free sugars are monosaccharides and disaccharides added to foods, sugars naturally present in honey, syrups,

fruit juices and fruit juice concentrates, and sugars from foods extracts and artificial flavourings, alcoholic drinks¹⁴.

RA reflect food typically consumed in one sitting and are developed and maintained mainly for nutrition labelling purposes. Nutrient intakes were computed using the Brazilian Food Composition Table. The Brazilian Geography and Statistic Institute provided the data from the RAs and estimation of the free sugars components were based on previous study methodology¹⁵.

Brazilian Dietary Index

Details on assignment of scores and alignment with the updated 2024 Pyramid food groups are provided in Table 1. Briefly, the index comprises 9 components named: (i) fruits and vegetables, (ii) (whole) grains, (iii) dairy, (iv) meats and eggs, (v) beans, (vi) fatty acids ratio (MONO + POLI/SAT), (vii) saturated fats, (viii) free sugars and (ix) sodium. Each component relates to key recommendations identified in the food pyramid groups. Scoring standards were taken directly from targets proposed in the pyramid groups for nutrients components or derived from the guidelines based on consumption data and expert judgment. Table 1 presented a list of the metric components, scores and standard scores. Fruits and vegetables (FV) and grains, with preference for whole grains should be the key component indicator for diet quality and healthy dietary patterns¹⁶. For example, FV and (whole) grains received the same points, but reference amount for each food differs based on their energy, fibre and grams¹⁰. Importantly, whole grains should be always given the preference, but we should not exclude other sources of grains to attend the context of each individual, not stigmatizing foods into “bad” and “good”¹⁶. A maximum score of 5 or 10 was given to components related to particular recommendation or closely linked recommendations. Then, all the 9 components were summed, and index has a maximum score of 100 and reflects adherence to healthy food choices of the 2024 Food Pyramid.

Table 1 - Brazilian Healthy Eating Index, 2024. Nutrition Focused Brazilian House Budget Survey, 2017-18.

Food Groups	Maximum points	Standard for maximum score	Standard for minimum score
Fruits and Vegetables	20	≥ 2.0 cup eq/1000 kcal	No fruits and vegetables
(Whole) grains [†]	20	≥ 128 grams eq/1000 kcal	No grains
Dairy	10	≥ 1.3 cup eq/1000 kcal	No dairy
Meat and Eggs	5	≥ 56 grams eq/1000 kcal	No meat and eggs
Beans	5	≥ 1.4 cup/1000 kcal	No beans
Fatty Acids ratio	10	(MUFA + PUFA)/SFA ≥ 2.5	(MUFA + PUFA)/SFA ≤ 1.2
Sodium	10	≤ 1.1g /1000kcal	≥ 2.0g/1000 kcal
Free sugars	10	≤ 6.5% of energy	≥ 26% of energy
Saturated fats	10	≤ 8% of energy	≥ 16% of energy
Total score	100	---	---

MUFA: Mono-unsaturated fats; PUFA: Poli-unsaturated fats; SFA: Saturated fats. [†]Preference for whole grains, such as brown rice, whole grain pasta, quinoa.

Statistical analysis

Six tasks were taken to evaluate the construct validity and reliability (internal consistency) of the dietary metric as predefined analyses of previous healthy eating indexes^{17,18}. The variability among individuals were assessed via distributions of usual dietary intakes for each components estimated using the National Cancer Institute multivariate Markov Chain Monte Carlo method¹⁹. Stratification by age and sex were conducted due within-individual random dietary intake variations that differ across life stages²⁰. A scoring algorithm was applied to the estimated usual intakes among pseudo-individuals to estimate total and components scores. The mean and percentiles of total dietary metric and components were estimated for total and sub samples. Mean dietary components and total scores for subgroups (age, sex, race and food security status) with expected differences in diet quality were estimated using the ratio method²¹. The hypothesis was tested with the U.S Healthy Eating Index, 2020 score² due to all the scores are intended to reflect overall diet quality. The relationship between the indexes were examined using Pearson correlation as well as linear regression with a restricted to account for potential non-linearity. Quantiles were selected a priori based on common placement to ensure enough data within each interval²². Association between energy intake and dietary metric scores was assessed using univariate linear regression and Pearson correlation. The last task for construct validity was the principal component analysis that was used to assess multidimensionality of the

metric based on the 9 components of the index and based on usual dietary intake. The eigenvalues and eigenvectors of the principal component analysis solution were assessed to confirm multidimensionality (variance of scores not explained by only one of its components). The association between component scores was also checked with Pearson correlation. Sampling weights were used to generalize results to the Brazilian population. Analyses were performed in R Studio Version 2023.06.1+524 (Posit Software, PBC).

RESULTS

Participants variability

Distribution of the total and components scores from the Brazilian metric based on usual dietary intake in Brazilians ≥ 10 yo are shown in Figure 1. The mean total score of the Brazilian index was of 65.65 (95%CI 65.55, 65.80) with a range from 40.11 (percentile 1) and 85.15 (percentile 99). Adolescents (64.36) and male adults (63.94) had the lowest total score and female adults (66.84) the highest. “Ceiling effects”, i.e., greater proportion of individuals receiving maximum score, were observed for grains, meats and eggs, beans, and free sugars.




Figure 1. Estimated distribution of total dietary scores based on usual dietary intakes in Brazilians ≥ 10 yo from the 2017-18 Nutrition Focus House Budget Survey, Brazil (n = 16,254).

Differences between components and total scores and subgroups

Mean differences of components scores from the Brazilian index with population subgroups are presented on Figure 2. Overall females vs. males and food secure vs. insecure presented better diet quality for fruits and vegetables and dairy. Adolescents vs. adults did not show significant differences between components.




Figure 2. Brazilian Healthy Eating Index – 2024 radar plots across the population subgroups, Nutrition focus House Budget Survey, 2017–18. *Each component score is plotted as mean of its maximum points on 9 different axes. The outer edge of the radar represents 20, 10 or 5 depending on the component, while the center represents 0 of the maximum score of any component. Food security is the lack of consistent, dependable access to sufficient food for an active and healthy life.*

Correlations between metrics and total energy intake

Table 2 depicts Pearson correlations between energy intake, components and total scores of the Brazilian metric. Sodium ($r = -0.52$) showed the strongest negative correlation with fruits and vegetables, while total score the strongest positive correlation with fatty acids ($r = 0.52$). Energy intake showed a stronger positive correlation for grains ($r = 0.71$) and negative correlation for fruits and vegetables ($r = -0.21$). There was a strong positive correlation between the Brazilian and the US index ($r = 0.75$) (data not shown).

Table 2. Pearson correlations between BHEI, 2024 components scores and energy intake of Brazilians ≥ 10 yo from the Nutrition focus House Budget Survey, 2017-18

	1	2	3	4	5	6	7	8	9	10	11
1	1.00	-0.03***	-0.02***	0.08***	-0.11***	0.05***	-0.03***	-0.52***	-0.09***	-0.11***	-0.21***
2		1.00	-0.01	0.08***	0.01	0.06***	-0.02***	0.03***	-0.32***	0.08***	0.71***
3			1.00	0.01*	0.03***	0.09***	0.01	-0.07***	-0.01**	0.10***	0.16***
4				1.00	-0.05***	-0.07***	-0.41***	-0.09***	-0.18***	-0.26***	0.23***
5					1.00	0.09***	0.07***	-0.01*	0.05***	0.03***	0.13***
6						1.00	0.13***	-0.11***	0.02**	0.12***	0.24***
7							1.00	0.02**	0.16***	0.52***	0.33***
8								1.00	0.09***	0.07***	0.31***
9									1.00	0.01*	0.00
10										1.00	0.43***
11											1.00

1 = Energy, 2 = C1 Fruits and vegetables; 3 = Grains, 4 = Dairy, 5 = Meats and eggs, 6 = Beans, 7 = Fatty acids ratio, 8 = Sodium, 9 = Free sugars, and 10 = Total Score. * p -value < 0.05; ** p -value < 0.01; *** p -value < 0.001

Principal components analysis of dietary components of the Brazilian Healthy Index

Scores from the first four principal components analysis (PCA) (i.e., linear combinations) of the Brazilian dietary metric components accounted for 60% of the total variance. Fatty acids, saturated fats and dairy were the 3 components the most to the 1st principal component accounting for the largest proportion of the total variance (21%). The other 4 PCAs explained further information on total scores patterns of the Brazilians consisting with the multidimensional nature of dietary intake. For example, fruits and vegetables, and free sugars presented the most variance for the 2nd PCA (15%); for the 3rd were grains and beans (13%) and the 4th grains and meat (11%) (Table 3).

DISCUSSION

The Brazilian Healthy Eating Index, 2024 was evaluated for construct validity and reliability to assess adherence to recommendation of the updated Food Pyramid, 2024¹⁰ on healthy food choices. The index showed sufficient variation across Brazilians with significant differences in population subgroups: sex and food insecurity status; had a strong correlation with the American Healthy Eating Index-2020 and captured several dimensions of diet quality. There was consistency between components correlations, e.g., grains and beans were positively correlated.

Table 3. Principal components analysis of components scores of Brazilian Healthy Eating Index, 2024 among Brazilians ≥ 10 yo. Nutrition focus House Budget Survey, 2017-2018.

	PC1	PC2	PC3	PC4
Fruits and Vegetables	-0.09	-0.62	0.31	-0.19
Grains	0.06	0.21	-0.44	0.36
Dairy	-0.49	-0.11	-0.11	0.01
Meats	0.13	-0.05	-0.31	-0.85
Beans	0.19	-0.30	-0.46	-0.13
Fatty acids	0.59	-0.09	0.08	0.08
Sodium	0.09	0.19	0.57	-0.25
Sugar	0.26	0.58	-0.20	-0.04
Saturated Fats	0.51	-0.29	0.16	0.16
% variance	21.46	15.19	13.06	10.97
% cumulative	21.46	36.65	49.70	60.67

The total diet score demonstrated variability among Brazilians ≥ 10 yo and by subgroups: adolescents, females ≥ 18 yo and males ≥ 18 yo. However, distributions of grains and meats and eggs were skewed for maximum scores and free sugars for minimum scores. In the case of meats and eggs and grains most individuals were meeting the recommendations, but in the case of free sugars were consuming above the recommendations (free sugars were reversed scored). Even though, variability of total Brazilian healthy eating index, 2024 scores was adequate to demonstrate differences in diet quality according to sex and food insecurity status². Further studies are needed to verify how other populations meet the recommendations of the updated 2024 food pyramid to be understood as good quality.

The strong correlation between the Brazilian healthy eating index, 2024 and the US Healthy Eating Index, 2020 reflected adherence to the 2020-2025 Dietary Guidelines for Americans². Construct validity was supported taking into account that both indices reflect adherence to sets of recommendations that are based on similar body of evidence¹⁰. In addition, variability among individuals for the Brazilian index showed similar variability with the US index demonstrating construct validity, i.e., 21.56 in percentile 1 and 66.89 in percentile 99.

The nine components of the Brazilian index were weighted differently to capture adequate proportions based on the food pyramid recommendations, similarly to American HEI, 2020² and the 2019 Canadian Food Guide²³. Most of the components were significantly correlated,

indicating overlap of dietary behaviours, leading to indirectly providing more weight to that dietary behaviour²⁴. The components free sugars ($r = 0.16$) and dairy ($r = -0.41$) were correlated with fatty acids, which can be explained by the fact that preparations or industrialized foods have both sugars and fats, and reduced consumption of dairy products or consumption low-fat dairies are linked to fatty acids. The effect of additional health and diet-relations should be investigated in future studies. Maybe weighting of the components can be change according to specific recommendations, e.g., specific population group (children vs. adults vs. older adults) or presence of disease or any other healthy condition that requires stricter cut-offs points. Nevertheless, modifying scoring standards may or may not reflect adherence to specific recommendations to the Food Pyramid as per the original purpose of the Brazilian index¹⁰. Further, females and food secure households, i.e., adequate access to food, showed a relatively better diet quality corroborating to other studies^{25,26}. Taken together, results suggest that intake of specific dietary components, i.e., fruits and vegetables, milk and dairy and sodium components, may be particularly sensitive to sex and food insecurity differences²⁷⁻²⁹.

The strengths of the present study were the use of population-based data consistent to the aim of the of the index and several metrics to evaluate psychometric properties of the Brazilian index. However, limitations should be noticed. First dietary assessment methods are prone to random and systematic errors that can reflect on the quality of the estimates of diet. The estimated mean of the two 24h-recalls are prone to these errors as compared to food frequency questionnaires³⁰. Second, changes over the time of the diet could not be assessed due to the cross-sectional design study. Third, evaluation among individuals with specific health conditions or other subgroups children and older people, could not be assessed. Fourth, Brazilian healthy eating index, 2024 was not assessed for health and diet health disparities. Lastly, the developing of a diet quality index based on the pyramid based on the Brazilian population might not allow for comparability to other countries and cultures.

CONCLUSIONS

The Brazilian Healthy Eating Index, 2024 showed good psychometric properties that reflects to the recommendations of healthy eating to the 2024 updated food pyramid. Future research of

Brazilian Healthy Eating Index, 2024 can be used as monitoring tool in public health research or as a tool for assessing a Brazilian dietary pattern and studying nutrition or diet health disparities.

REFERENCIAS

- (1) English LK, Ard JD, Bailey RL, Bates M, Bazzano LA, Boushey CJ, Brown C, Butera G, Callahan EH, Jesus J de, Mattes RD, Mayer-Davis EJ, Novotny R, Obbagy JE, Rahavi EB, Sabate J, Snetselaar LG, Stookey EE, Horn LV, Venkatramanan S, Heymsfield SB. Evaluation of Dietary Patterns and All-Cause Mortality: A Systematic Review. *JAMA Network Open*. 2021;4(8):E2122277. <https://doi.org/10.1001/jamanetworkopen.2021.22277>
- (2) Shams-White MM, Pannucci TE, Lerman JL, Herrick KA, Zimmer M, Mathieu KM, Stookey EE, Reedy J. Healthy Eating Index-2020: Review and Update Process to Reflect the Dietary Guidelines for Americans, 2020-2025. *J Acad Nutr Diet*. 2023;123(9):1280–1288. <https://doi.org/10.1016/j.jand.2023.05.015>
- (3) Woodside J, Young IS, McKinley MC. Culturally adapting the Mediterranean Diet pattern - a way of promoting more “sustainable” dietary change? *Br J Nutr*. 2022;128(4):693–703. <https://doi.org/10.1017/S0007114522001945>
- (4) Aljuraiban GS, Gibson R, Oude Griep LM, Okuda N, Steffen LM, Van Horn L, Chan Q. Perspective: The Application of A Priori Diet Quality Scores to Cardiovascular Disease Risk—A Critical Evaluation of Current Scoring Systems. *Adv Nutr*. 2020;11(1):10–24. <https://doi.org/10.1093/advances/nmz059>
- (5) Dalwood P, Marshall S, Burrows TL, McIntosh A, Collins CE. Diet quality indices and their associations with health-related outcomes in children and adolescents: an updated systematic review. *Nutr J*. 2020;19(1):118. <https://doi.org/10.1186/s12937-020-00632-x>
- (6) Bromage S, Batis C, Bhupathiraju SN, Fawzi WW, Fung TT, Li Y, Deitchler M, Angulo E, Birk N, Castellanos-Gutiérrez A, He Y, Fang Y, Matsuzaki M, Zhang Y, Moursi M, Gicevic S, Holmes MD, Isanaka S, Kinra S, Sachs SE, Stampfer MJ, Stern D, Willett WC. Development and Validation of a Novel Food-Based Global Diet Quality Score (GDQS). *J Nutr*. 2021;151(12 Suppl 2):75S-92S. <https://doi.org/10.1093/jn/nxab244>
- (7) Norde MM, Bromage S, Marchionni DML, Vasques AC, Deitchler M, Arsenault J, de Carvalho AM, Velloso L, Willett W, Giovannucci E, Geloneze B. The global diet quality score as an indicator of adequate nutrient intake and dietary quality – a nation-wide representative study. *Nutr J*. 2024;23(1):42. <https://doi.org/10.1186/s12937-024-00949-x>
- (8) Agurs-Collins T, Alvidrez J, ElShourbagy Ferreira S, Evans M, Gibbs K, Kowtha B, Pratt C, Reedy J, Shams-White M, Brown AG. Perspective: Nutrition Health Disparities Framework: A Model to Advance Health Equity. *Advances in Nutrition*. 2024;15(4):100194. <https://doi.org/10.1016/j.advnut.2024.100194>
- (9) FAO, WHO. Food security indicators – latest updates and progress towards ending hunger and ensuring food security [Internet]. FAO/OMS; 2023. Available from: <https://www.fao.org/3/cc3017en/online/state-food-security-and-nutrition-2023/food-security-nutrition-indicators.html>

- (10) Philippi ST, Aquino R de C, da Silva EAFT, Leme ACB. Healthy lifestyle and sustainable behaviours: a proposal to the food pyramid for the Brazilian context. *Nutrition*. 2025;112771. <https://doi.org/10.1016/j.nut.2025.112771>
- (11) Men F, Tarasuk V. Classification Differences in Food Insecurity Measures between the United States and Canada: Practical Implications for Trend Monitoring and Health Research. *J Nutr*. 2022;152(4):1082–1090. <https://doi.org/10.1093/jn/nxab447>
- (12) Fisberg RM, Slater B, Barros RR, Lima FD de, Cesar CLG, Carandina L, Barros MB de A, Goldbaum M. Índice de Qualidade da Dieta: avaliação da adaptação e aplicabilidade. *Revista de Nutriçãor*. 2004;17:301–318. <https://doi.org/10.1590/S1415-52732004000300003>
- (13) Giavarina D. Understanding Bland Altman analysis. *Biochem Med (Zagreb)*. 2015;25(2):141–151. <https://doi.org/10.11613/BM.2015.015>
- (14) Swan GE, Powell NA, Knowles BL, Bush MT, Levy LB. A definition of free sugars for the UK. *Public Health Nutr*. 2018;21(9):1636–1638. doi: 10.1017/S136898001800085X.
- (15) Rana H, Mallet MC, Gonzalez A, Verreault MF, St-Pierre S. Free Sugars Consumption in Canada. *Nutrients*. 2021 Apr 27;13(5):1471. <https://doi.org/10.1017/S136898001800085X>
- (16) World Health Organization. Healthy diet [Internet]. 2020 [cited 2024 Feb 9]. Available from: <https://www.who.int/news-room/fact-sheets/detail/healthy-diet>
- (17) Krebs-Smith SM, Pannucci TE, Subar AF, Kirkpatrick SI, Lerman JL, Tooze JA, Wilson MM, Reedy J. Update of the Healthy Eating Index: HEI-2015. *J Acad Nutr Diet*. 2018;118(9):1591–1602. <https://doi.org/10.1016/j.jand.2018.05.021>
- (18) Guenther PM, Reedy J, Krebs-Smith SM. Development of the Healthy Eating Index-2005. *J Am Diet Assoc*. 2008;108(11):1896–1901. <https://doi.org/10.1016/j.jada.2008.08.016>
- (19) Zhang S, Midthune D, Guenther PM, Krebs-Smith SM, Kipnis V, Dodd KW, Buckman DW, Tooze JA, Freedman L, Carroll RJ. A new multivariate measurement error model with zero-inflated dietary data and its application to dietary assessment. *Ann Appl Stat*. 2011;5(2B):1456–1487. <https://doi.org/10.1214/10-AOAS446>
- (20) Herrick KA, Rossen LM, Parsons R, Dodd KW. Estimating Usual Dietary In take From National Health and Nutrition Examination Survey Data Using the National Cancer Institute Method. *Vital Health Stat 2*. 2018;(178):1–63.
- (21) Freedman LS, Commins JM, Moler JE, Arab L, Baer DJ, Kipnis V, Midthune D, Moshfegh AJ, Neuhaus ML, Prentice RL, Schatzkin A, Spiegelman D, Subar AF, Tinker LF, Willett W. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake. *Am J Epidemiol*. 2014;180(2):172–188. <https://doi.org/10.1093/aje/kwu116>
- (22) Harrell FE. General Aspects of Fitting Regression Models. In: Harrell Jr Frank E, editor. *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis* [Internet]. Cham: Springer International Publishing; 2015 [cited 2024 Dec 26]. p. 13–44. Available from: https://doi.org/10.1007/978-3-319-19425-7_2
- (23) Brassard D, Elvidge Munene LA, St-Pierre S, Gonzalez A, Guenther PM, Jessri M, Vena J, Olstad DL, Vatanparast H, Prowse R, Lemieux S, L'Abbe MR, Garriguet D, Kirkpatrick SI, Lamarche B. Evaluation of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices. *Appl Physiol Nutr Metab*. 2022;47(5):582–594. <https://doi.org/10.1139/apnm-2021-0416>

- (24) van Lee L, Geelen A, van Huysduynen EJCH, de Vries JHM, van't Veer P, Feskens EJM. The Dutch Healthy Diet index (DHD-index): an instrument to measure adherence to the Dutch Guidelines for a Healthy Diet. *Nutr J*. 2012;11(1):49. <https://doi.org/10.1186/1475-2891-11-49>
- (25) Leme A, Muszynski D, Mirotta JA, Carroll N, Hogan JL, Jewell K, Yu J, Fisberg RM, Duncan AM, Ma DWL, Haines J, Guelph Family Health Study null. Diet Quality of Canadian Preschool Children: Associations with Socio-demographic Characteristics. *Can J Diet Pract Res*. 2021;82(3):131–135. <https://doi.org/10.3148/cjdpr-2021-009>
- (26) McNaughton SA. Dietary Patterns and Diet Quality: Approaches to Assessing Complex Exposures in Nutrition. *Australasian Epidemiologist*. 2020;17(1):35–37.
- (27) Dalwood P, Marshall S, Burrows TL, McIntosh A, Collins CE. Diet quality indices and their associations with health-related outcomes in children and adolescents: an updated systematic review. *Nutr J*. 2020;19(1):118. <https://doi.org/10.1186/s12937-020-00632-x>
- (28) Kang M, Park SY, Shvetsov YB, Wilkens LR, Marchand LL, Boushey CJ, Paik HY. Sex differences in sociodemographic and lifestyle factors associated with diet quality in a multiethnic population. *Nutrition*. 2019;66:147–152. <https://doi.org/10.1016/j.nut.2018.11.022>
- (29) Thomson JL, Tussing-Humphreys LM, Goodman MH, Landry AS. Diet quality in a nationally representative sample of American children by sociodemographic characteristics. *Am J Clin Nutr*. 2019;109(1):127–138. <https://doi.org/10.1093/ajcn/nqy284>
- (30) Kirkpatrick SI, Baranowski T, Subar AF, Tooze JA, Frongillo EA. Best Practices for Conducting and Interpreting Studies to Validate Self-Report Dietary Assessment Methods. *J Acad Nutr Diet*. 2019;119(11):1801–1816. <https://doi.org/10.1016/j.jand.2019.06.010>

AUTHORS' CONTRIBUTIONS

ACBL and STP was involved in the conception and design of the study and interpretation of analysis. ACBL conducted statistical analysis and wrote the first draft of this manuscript. STP was involved in the adaption of the food pyramid to the Brazilian context, to its dietary recommendations and to critically review this manuscript. All authors approved the final version of this manuscript.

FUNDING

The authors declare that there has been no funding to carry out this study.

CONFLICTS OF INTEREST

The authors state that there are no conflicts of interest when writing the manuscript.

DATA AVAILABILITY

This study was a secondary analysis of the Brazilian House Budget Survey – 2017-18 data. All information was de-identified and vetted by a trained member of the Brazilian Geography Statistics Institute prior to their publicly released. Data can be found:

<https://www.ibge.gov.br/estatisticas/sociais/saude/24786-pesquisa-de-orcamentos-familiares-2.html?=&t=microdados>